

DETERMINISTIC CHAOS IN MATERIALS EXHIBITING PHASE
TRANSITIONS(U) MINNESOTA UNIV MINNEAPOLIS M SLEMMOD
01 JUN 83 AFOSR-TR-83-0929 AFOSR-82-0246

UNCLASSIFIED

F/G 20/13


NL

END

DATE _____

VALUED

28



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

(2)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 88-0929	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) "FINAL REPORT-AFOSR-82-0246"		5. TYPE OF REPORT & PERIOD COVERED FINAL
7. AUTHOR(s) Marshall Slemrod		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Minnesota Minneapolis MN 55455		8. CONTRACT OR GRANT NUMBER(s) AFOSR-82-0246
11. CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NM Bldg. 410 Bolling AFB DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F; 2304/A4
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1, 1983
		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) M. Slemrod spent one half of the Spring 1983 semester at the Institute for Mathematics and its Applications. During that time he interacted with colleagues, engaged in research, and gave two public lectures at the Institute: (i) Chaos in Phase Transitions, (ii) Dynamics of Phase Transitions. The main thrust of Slemrod's research was in two areas, specifically: (i) Deterministic chaos in materials exhibiting phase transitions, and cont. (over)		

DTIC
ELECTE
S NOV 29 1983 D
A E

DD FORM 1 JAN 73 1473

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD-A135172

DTIC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

(ii) Admissibility criteria for weak solutions of the non-hyperbolic conversation laws which describe dynamic phase transitions.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

AFOSR-TR- 83 - 0929

Final Progress Report

**Marshall Slemrod
Institute for Mathematics and its Applications
University of Minnesota
Minneapolis, Minnesota 55455**

1 June 1983

AFOSR - 82-0246

**Approved for public release;
distribution unlimited.**

83 11 29 272

0. Introduction

→ ~~The author~~
~~M. Slater~~ spent one half of the Spring 1983 semester at the Institute for Mathematics and its Applications. During that time he interacted with colleagues, engaged in research, and gave two public lectures at the Institute:

~~(i) Chaos in Phase Transitions, and~~

~~(ii) Dynamics of Phase Transitions.~~

^{his}
The main thrust of ~~Slater's~~ research was in two areas, specifically:

~~(i) Deterministic chaos in materials exhibiting phase transitions, and~~

~~(ii) Admissibility criteria for weak solutions of the non-hyperbolic conservation laws which describe dynamic phase transitions.~~


A detailed description of this research is given below.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for release to IAW AFR 197-12.
Distribution is unlimited.
MATTHEW J. KEMPER
Chief, Technical Information Division

1. Chaos in Materials Exhibiting Phase Transitions

In recent work Holmes and Marsden (1981) have employed the Mel'nikov (1963) technique to establish the existence of deterministic chaos in periodically forced evolution equations. In this work Slemrod (in collaboration with J. E. Marsden of the University of California, Berkeley) combined this program by analyzing two problems arising from the vander Waals theory of phase transitions [see vander Waals (1893, 1979)].

The first problem considered was dynamic spinoidal decomposition. Assume we have a vander Waals fluid with isotherms as shown in Figure 1.



Here p denotes the pressure given by the vander Waals constitutive relation

$$p(w, \theta) = \frac{R\theta}{w-b} - \frac{a}{w^2} \quad (1.1)$$

R , b , a are positive constants, ρ is the density, $w = 1/\rho$ the specific volume, and θ is the absolute temperature. Figure 1 sketches the vander Waals isotherms

for θ above, equal to, and below the critical temperature $\theta_{\text{crit}} = 8a/27bR$.
 [A good reference is Fermi (1936).] For $\theta_0 < \theta_{\text{crit}}$, $p(w, \theta)$ has the following features:

- (i) $p_w(w, \theta_0) < 0$ on $(b, \alpha) \cup (\beta, \infty)$,
- (ii) $p_w(w, \theta_0) > 0$ on (α, β) ,
- (iii) $p_w(\alpha, \theta_0) = p_w(\beta, \theta_0) = 0$.

The domain (b, α) corresponds to the fluid being liquid; the domain (β, ∞) corresponds to the fluid being vapor; the domain (α, β) is the unstable region and is referred to as the spinodal. Also the point w_0 is where $p_w = 0$ on the graph of the θ_0 isotherm (w_0 is the zero of $R\theta_0 w^4 + a(w-b)^3 = 0$).

The fluid flow is thought of as taking place along the x -axis in a tube of unit cross section of fixed volume. If $\theta_1 > \theta_{\text{crit}}$ the state $w = w_0$ will describe a stable homogeneous configuration for the hydrodynamic equations [see Felderhof (1970), Slemrod (1982)]. We then "instantaneously" quench the fluid by reducing the temperature to $\theta_0 < \theta_{\text{crit}}$. The homogeneous state w_0 will now be in the unstable spinodal region of the θ_0 isotherm. We studied the effect of a small periodic in time fluctuation of the absolute temperature about θ_0 . Specifically we showed how the loss of stability of w_0 is accompanied by deterministic chaos in that there will be a Poincaré-Birkhoff-Smale horseshoe in the dynamics.

The second problem considered is the effect of a small thermal perturbation, periodic in space, of the form $\theta(x) = \theta_0 + \epsilon \cos qx$, ϵ small, on the equilibrium configuration of an infinite tube of liquid under given applied load. In this case we show there are solutions with the features of both metastable and co-existing phases that exhibit spatial chaos.

It seems interesting to note that the above equilibrium result has features in common with experimental observations presented by Dr. L. Zapas of NBS at the IMA Workshop on Orienting Polymers, March 21-25, 1983.

The results of this research will appear as an IMA report and will be submitted for publication.

2. Admissibility Criteria for Weak Solutions of Conservation Laws

In recent years an ever-increasing number of people have taken up Gelfand's (1963) program of parabolic regularization of quasilinear conservation laws. While this work has usually been motivated by the search for reasonable admissibility criteria for weak solutions of the conservation laws, it has also provided a technique for proving existence of solutions as well. Specifically in this regard we note the work of Oleinik (1957) for a scalar conservation law and the recent remarkable results of DiPerna (1983) for a two-dimensional system of conservation laws. In light of this work Slemrod attempted to give (where possible) a physical motivation to the Gelfand program with respect to the equations of compressible fluid flow.

Consider a system of n conservation laws written in vector form

$$F(U)_t + G(U)_x = 0 \quad (2.1)$$

where $U(x, t)$ is an n -vector, $-\infty < x < \infty$, and $t > 0$. As is well known, the initial value problem (2.1) and

$$U(x, 0) = U_0(x) \quad (2.2)$$

does not in general possess global smooth solutions if (2.1) is nonlinear, no matter how smooth U_0 . Hence we are led to search for weak solutions of (2.1). Unfortunately, there is a serious price to pay for enlarging the solution class, namely loss of uniqueness of solutions.

The Gelfand program of parabolic regularization attempts to give a way to pick out the physically relevant weak solutions of (2.1) and thereby, hopefully, recover uniqueness of weak solutions of (2.1). Specifically Gelfand suggested that the physically admissible solutions of (2.1) should be limits of solution U^ϵ of the "artificially viscous" equations

$$F(U^\epsilon)_t + G(U^\epsilon)_x = \epsilon D(U^\epsilon)_{xx} \quad (2.3)$$

where $D: \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\epsilon > 0$ a small parameter. For example, the equations of viscous isothermal compressible fluid flow in Lagrange coordinates are

$$u_t^\epsilon + p(w^\epsilon)_x = \epsilon u_{xx}^\epsilon$$

(2.4)

$$w_t^\epsilon - u_x^\epsilon = 0$$

so that $U^\epsilon = \begin{pmatrix} u^\epsilon \\ w^\epsilon \end{pmatrix}$ and $D = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$.

Unfortunately D in this example is singular, whereas the recent results of DiPerna on existence of weak solutions of (2.1) as limits of (2.3) require D nonsingular.

Slemrod in his research attempted to give a physical justification for nonsingular D matrices based on the vander Waals-Korteweg gradient theory of stress. Using this idea Slemrod was able to apply DiPerna's technique to prove the existence of weak solutions of (2.4, $\epsilon = 0$) for all $t \geq 0$ when p is described by the θ_{crit} isotherm of Fig. 1.

This work will appear as an IMA report and will be submitted for publication.

References

- DiPerna, R. (1983), Convergence of approximate solutions to conservation laws, to appear in *Archive for Rational Mechanics and Analysis*.
- Felderhof, B.U. (1970), Dynamics of the diffuse gas-liquid interface near the critical point, *Physica*, 48, 541-560.
- Fermi, E. (1936), Thermodynamics, New York: Dover.
- Gelfand, I. (1963), English translation of "Some problems in the theory of quasi-linear equations," *Amer. Math. Soc. Transl. Ser. 2*, 6, 295-381.
- Holmes, P. and J.E. Marsden (1981), A partial differential equation with infinitely many periodic orbits: chaotic oscillations of a forced beam, *Archive for Rational Mechanics and Analysis*, 76, 135-166.
- Mel'nikov, V.K. (1963), On the stability of the center for time periodic perturbations, *Trans. Moscow Math. Society*, 12, 1-57.
- Oleinik, O. (1957), English translation of "Discontinuous solutions of nonlinear differential equations," *Amer. Math. Soc. Transl. Ser. 2*, 26, 95-172.
- Slemrod, M. (1982), Admissibility criteria for propagating phase boundaries in a vander Waals fluid, *Archive for Rational Mechanics and Analysis*, 81, 301-315.
- vander Waals, J.D. (1893), *Venhandel, Konink. Akad. Wetenschapen. Amsterdam*, Vol. 1, No. 8; (1979), Translation of J.D. vander Waals "The thermodynamic theory of capillarity under the hypothesis of a continuous variation of density," *J. Statistical Physics*, 20, 197-244.